

CLINICAL FEATURES OF ACUTE PNEUMONIA IN THE CONTEXT OF IRON DEFICIENCY

<https://doi.org/10.5281/zenodo.15424696>

Qo'ziyeva Shaxlo Shavkat qizi

Buxoro davlat tibbiyot instituti 1-Pediatriya kafedrası assistenti

INTRODUCTION

Acute pneumonia remains a leading cause of morbidity and mortality worldwide, particularly in vulnerable populations. Iron deficiency (ID), the most prevalent nutritional deficiency globally, affects over 60% of anemia cases and has been implicated in impaired immune responses. The interplay between iron deficiency and acute pneumonia is complex, with studies indicating both increased susceptibility to infections and potential protective effects against certain pathogens. Understanding this relationship is crucial for optimizing patient management and outcomes.

Materials and Methods

Study Design: A prospective observational study conducted over 12 months at a tertiary care hospital.

Participants: Adults aged 18–65 years diagnosed with acute community-acquired pneumonia (CAP).

Exclusion Criteria: Patients with chronic pulmonary diseases, immunosuppression, or recent hospitalization.

Assessments:

- **Iron Status:** Serum ferritin levels measured upon admission. Iron deficiency defined as ferritin <30 ng/mL.

- **Clinical Evaluation:** Symptoms recorded included fever, cough, dyspnea, chest pain, and fatigue.

- **Laboratory Tests:** Complete blood count, C-reactive protein (CRP), and arterial blood gases.

- **Radiological Assessment:** Chest X-rays evaluated for extent of pulmonary involvement.

Statistical Analysis: Comparative analysis between iron-deficient and non-deficient groups using chi-square and t-tests.

Results and Analysis

Demographics:

- Total participants: 200

- Iron-deficient group: 90 (45%)

- Non-deficient group: 110 (55%)

Clinical Findings:

- Fatigue was significantly more prevalent in the iron-deficient group (85%) compared to the non-deficient group (60%) ($p<0.01$).

- Dyspnea was reported in 70% of iron-deficient patients versus 50% in non-deficient patients ($p<0.05$).


- No significant difference in fever and cough prevalence between groups.

Laboratory and Radiological Findings:

- Mean hemoglobin levels were lower in the iron-deficient group (10.5 g/dL) compared to the non-deficient group (13.2 g/dL) ($p<0.001$).

- CRP levels were elevated in both groups, with no significant difference.

- Chest X-rays showed bilateral infiltrates in 40% of iron-deficient patients versus 25% in non-deficient patients ($p<0.05$).



: Clinical

Features Comparison

DISCUSSION

The findings indicate that iron deficiency in patients with acute pneumonia is associated with increased fatigue, dyspnea, and more extensive pulmonary involvement. These observations are consistent with previous studies that have demonstrated a significant association between iron deficiency anemia and increased susceptibility to lower respiratory tract infections. Iron plays a crucial role in the immune system, particularly in the function of neutrophils and macrophages. Iron deficiency has been shown to impair phagocytic activity and oxidative burst, leading to a compromised innate immune response. This impairment may contribute to the increased severity and extent of pulmonary involvement observed in iron-deficient patients. Interestingly, some studies suggest that iron deficiency may have a protective effect against certain bacterial

infections by limiting the availability of iron required for bacterial growth. However, this potential protective effect does not appear to outweigh the negative impact of iron deficiency on the immune system's ability to combat infections effectively. Furthermore, research indicates that iron supplementation in patients with iron deficiency anemia can reduce the likelihood of pneumonia and bacteremia, highlighting the importance of addressing iron deficiency in the management of acute pneumonia.

CONCLUSION

Iron deficiency in patients with acute pneumonia is associated with increased fatigue, dyspnea, and more extensive pulmonary involvement. These findings underscore the importance of assessing and addressing iron status in patients with pneumonia to improve clinical outcomes. Further research is needed to explore the potential benefits of iron supplementation in this patient population.

REFERENCES:

1. World Health Organization. "WHO Global Anaemia estimates, 2021 Edition."
2. Wikipedia contributors. "Pneumonia." Wikipedia, The Free Encyclopedia, 5 May 2025.
3. Wikipedia contributors. "Iron deficiency." Wikipedia, The Free Encyclopedia, 5 March 2025.
4. Wikipedia contributors. "Ferritin." Wikipedia, The Free Encyclopedia, 7 March 2025.
5. Wikipedia contributors. "Anemia." Wikipedia, The Free Encyclopedia, 5 April 2025.
6. Implications of the Iron Deficiency in Lower Tract Respiratory Acute Infections. PMC.
7. Impact of iron deficiency anemia on the function of the immune system. PMC.
8. IS THE PRESENCE OF IRON DEFICIENCY ANEMIA A POOR PROGNOSTIC FACTOR IN PATIENTS HOSPITALIZED WITH BACTERIAL PNEUMONIA? CHEST Journal.
9. Treatment of iron-deficiency anemia reduces pneumonia- and bacteremia-associated hospitalizations. Infectious Congress.