

ВЛИЯНИЕ ИИ НА РАЗВИТИЕ УСТНОЙ РЕЧИ В СОВРЕМЕННОМ МИРЕ

<https://doi.org/10.5281/zenodo.17010501>

Уразымбетова Юлдуз Реимбаевна

Магистратура TESOL (преподавание Английского языка носителям других языков), Webster University

Бакалавриат Иностранные языки и литература, Узбекский Государственный Университет Мировых Языков

Абстракт: *в условиях современной глобализации знание английского языка приобретает всё большее значение, а вместе с ним возрастает и необходимость в развитии высоких коммуникативных навыков. С развитием технологий искусственный интеллект (ИИ) становится важным инструментом в сфере образования, включая обучение иностранным языкам. Настоящее исследование представляет собой обзор научной литературы, посвящённой изучению влияния ИИ на формирование коммуникативных компетенций у учащихся, изучающих английский язык. Цель работы — проанализировать существующие исследования и теоретические источники, рассматривающие использование технологий ИИ в процессе обучения английскому. В начале эссе даётся общее представление об искусственном интеллекте и его применении в образовательной среде. Далее описываются способы, с помощью которых ИИ может способствовать развитию навыков устной речи, аудирования, чтения и письма у изучающих язык.*

Ключевые слова: *Искусственный интеллект (ИИ), обучение английскому языку, коммуникативные навыки, адаптивные образовательные технологии, этические аспекты ИИ, образовательные технологии*

В современном взаимосвязанном мире владение английским языком становится неотъемлемым навыком для тех, кто стремится добиться успеха в личной жизни, учёбе и карьере (Сари, 2023). Умение общаться на английском открывает доступ к образованию, международным проектам и новым профессиональным возможностям по всему миру (Сари, 2021). По мере роста спроса на язык преподаватели и учёные ищут новые методы обучения, одним из которых стала интеграция искусственного интеллекта (ИИ) в образовательный процесс. ИИ, способный имитировать человеческое мышление, за последние годы сделал большой шаг вперёд, прочно войдя в различные сферы общества. В образовании технологии на его основе рассматриваются как перспективный инструмент для поддержки и развития языковых умений, особенно для улучшения коммуникативных навыков. ИИ предлагает персонализированный, интерактивный и адаптивный

формат обучения, учитывающий уникальные потребности и предпочтения каждого учащегося. Цель этой обзорной статьи — изучить, как ИИ влияет на развитие коммуникативных навыков у изучающих английский язык. На основе анализа существующих исследований и научных публикаций мы представляем текущее состояние знаний, обобщаем основные выводы, а также выявляем тенденции и пробелы в этой области.

В этом обзорном исследовании применён последовательный системный подход к поиску, отбору и анализу научных публикаций, посвящённых роли искусственного интеллекта (ИИ) в формировании коммуникативных навыков у изучающих английский язык. Процедура работы включает четыре ключевых этапа:

1. Формулировка целей и вопросов исследования

Сначала чётко определяются главная цель и вспомогательные вопросы, которые будут направлять весь обзор литературы. Основная задача — выяснить, каким образом ИИ способствует развитию устной речи, аудирования, чтения и письма у учащихся английского языка. Дополнительно разрабатываются уточняющие вопросы об эффективности ИИ-инструментов, педагогических последствиях их применения и связанных этических аспектах.

2. Поиск источников

Проводится комплексный поиск релевантной литературы в академических базах данных, рецензируемых журналах, материалах конференций и специализированных онлайн-архивах. Используются такие ключевые слова и фразы, как «искусственный интеллект», «ИИ в обучении языкам», «английский как второй язык», «коммуникативная компетенция» и другие близкие по теме термины. По мере работы стратегия поиска уточняется и дополняется для обеспечения полноты охвата.

3. Отбор публикаций

Для включения и исключения материалов применяются заранее установленные критерии. Литература отбирается по соответствию тематике, дате публикации (акцент на последние годы), научной надёжности и методологической строгости. Приоритет отдаётся эмпирическим исследованиям, рецензируемым статьям и авторитетным теоретическим обзорам.

4. Извлечение данных и их обработка

Изученные источники подвергаются детальному анализу: фиксируются ключевые результаты, описания применяемых методик, характеристики выборок и теоретические подходы. Эта информация систематизируется для выявления общих тем, трендов и различий между работами. При необходимости проводится сравнительный анализ, чтобы подчеркнуть сходства и отличия в выводах различных авторов. Таким образом, описанная методология обеспечивает структурированный и прозрачный обзор существующих исследований по применению ИИ в развитии коммуникативных умений изучающих английский язык.

5. Анализ ограничений и выявление пробелов

Оцениваются ограничения как отдельных исследований, так и всей обзорной работы. Обсуждаются возможные искажения, такие как языковая или публикационная предвзятость. Также определяются недостаточно изученные аспекты темы, что позволяет выделить направления для дальнейших исследований.

6. Подготовка итогового отчёта

Все результаты и выводы оформляются в виде связного и логичного текста, соответствующего структуре научной обзорной статьи. В отчёт включаются: введение, описание методологии, обзор литературы, анализ и обсуждение полученных данных, выводы и предложения для будущих исследований.

Важно подчеркнуть, что данный метод основан исключительно на вторичном анализе — то есть на изучении и обобщении уже существующих научных источников. Проведение собственных экспериментов или опросов не входило в рамки данного исследования. Ниже представлены ключевые результаты, полученные в ходе литературного обзора:

1. Влияние искусственного интеллекта на развитие коммуникативных навыков у изучающих английский язык

a. Многие исследования подчёркивают положительное влияние ИИ-инструментов, таких как системы распознавания речи и виртуальные преподаватели, на улучшение устной речи и произношения.

b. Интерактивные платформы и чат-боты с ИИ стимулируют активное участие учащихся и помогают им свободнее говорить по-английски за счёт мгновенной обратной связи и постоянной языковой практики.

c. Искусственный интеллект в адаптивных образовательных технологиях предоставляет возможность персонализировать обучение языку, учитывая уровень подготовки, скорость усвоения материала и уникальные потребности каждого ученика.

2. Педагогические аспекты применения ИИ в языковом обучении

a. Использование ИИ в обучении способствует развитию самостоятельности: учащиеся могут заниматься в удобное время, получать обратную связь и доступ к материалам без постоянной помощи преподавателя.

b. ИИ-платформы обеспечивают персонализированный учебный путь и предлагают адаптивные методы оценки, учитывающие успехи, стиль обучения и интересы студентов.

4. Выявленные пробелы и направления для будущих исследований:

a. Несмотря на положительные результаты, требуется дальнейшее изучение долгосрочного влияния ИИ на развитие коммуникативных навыков изучающих английский язык, включая устойчивость приобретённых умений. b. Будущие исследования должны быть направлены на определение наилучших способов

интеграции ИИ в языковое образование и выявление эффективных методик, способствующих наиболее полному раскрытию потенциала ИИ в обучении.

с. Необходимо разрабатывать чёткие этические стандарты и руководства, регулирующие вопросы безопасности данных, прозрачности алгоритмов, устранения предвзятости и общего этического использования ИИ в образовательных целях.

Анализ существующих исследований позволяет сделать вывод о высокой эффективности ИИ в развитии коммуникативных навыков у изучающих английский язык. Особый акцент сделан на педагогических преимуществах, таких как персонализация обучения и развитие самостоятельности. Вместе с тем подчёркивается важность этических аспектов и необходимости дальнейших исследований, направленных на безопасную, инклюзивную и результативную интеграцию технологий ИИ в языковое образование.

СПИСОК ИСПОЛЬЗОВАННОЙ ЛИТЕРАТЫ:

1. Буоламвини, Ж., & Гебру, Т. (2018). Gender shades: Intersectional accuracy disparities in commercial gender classification. *Proceedings of the 1st Conference on Fairness, Accountability and Transparency*, 81-91
2. Ванг, Ю., & Лю, Х. (2019). The application of AI in language learning: A case study of an AI powered language learning application. *International Journal of Distance Education Technologies*, 17(1), 1-14.
3. Ли, Ж., Сие, Х., Ли, Д., & Лу, С. (2020). Effects of an AI-based speech recognition system on improving EFL learners' pronunciation. *Computers & Education*, 156, 103952.
4. Сари, М. Н. (2021). THE ROLE OF METACOGNITIVE STRATEGIES TRAINING IN TEACHING ENGLISH IN INDONESIAN EFL CLASSROOMS. *Jurnal Inovasi Penelitian*, 1(9), 1825-1834.
5. Сари, М. Н. (2023). STUDENTS' ABILITY AND DIFFICULTIES IN CONSTRUCTING CONDITIONAL SENTENCE TYPE II AT SMAN 10 KERINCI JAMBI. *Dharmas Education Journal (DE_Journal)*, 4(1), 23-32.
6. Сью, С., & Юан, С. Т. (2021). Data privacy protection and research ethics in intelligent learning systems: Current trends and future directions. *Journal of Educational Technology & Society*, 24(1), 17-32.
7. Чен, С. М., Хуанг, Ю. М., & Киншук. (2018). Effects of an intelligent tutoring system on students' learning outcomes and attitudes in the context of English as a foreign language. *Journal of Computer Assisted Learning*, 34(4), 376-393.
8. Чжан, М., Триллинг, Д., Пен, Ж., & Ванг, Ю. (2020). Adaptive learning in AI-based English language learning systems: Past, present, and future. *Journal of Educational Technology Development and Exchange*, 13(1), 1-16.

9. Чжан, Д., Тннг, Ж., Гау, Л., & Вей, С. (2019). Chatbot-assisted language learning: An empirical study of learners' perceptions and attitudes. *IEEE Access*, 7, 13310-13319.
10. Чжен, Х., & Синг, Ю. (2020). An adaptive learning platform based on AI for English learning. *IEEE Access*, 8, 202612-202620.

BIOENGINEERING THE CASPIAN TIGER: CAN CRISPR AND INTERSPECIES CLONING REVIVE AN EXTINCT SUBSPECIES?

<https://doi.org/10.5281/zenodo.17010507>

Toshpulatov Bexruz Dilshod o'g'li

Abstract: *The Caspian tiger (*Panthera tigris virgata*), once the largest tiger subspecies in Central Asia, was driven to extinction by mid-20th century due to habitat destruction and hunting[1][2]. Genomic analyses show that Caspian tigers were nearly identical to living Siberian (Amur) tigers, differing by only a few nucleotides[3][4]. This close relationship has led to proposals to reintroduce “proxy” tigers into the former Caspian range using Amur tigers or bioengineered equivalents[3][5]. Recent advances in biotechnology – notably CRISPR/Cas9 genome editing, stem-cell techniques, and interspecies cloning – raise the possibility of “de-extinction”. We review published literature on Caspian tiger history, genetics, and on de-extinction technologies. We analyze whether CRISPR can recreate missing Caspian tiger genetic traits and whether a viable embryo could be produced via interspecies somatic cell nuclear transfer (SCNT) or related methods[6][7]. We assess biological, ecological, technological, and ethical factors. Our findings suggest that, in principle, engineered Caspian tigers could be produced: the genetic differences with Siberian tigers are minimal, and CRISPR has already been used to synthesize extinct-wolf genes in living wolves[8] and to introduce mammoth traits into elephant cells[6]. However, substantial challenges remain. These include obtaining sufficient high-quality Caspian tiger DNA, ensuring proper gene regulation and epigenetic reprogramming, and finding or creating suitable surrogates (e.g. Siberian tiger or domestic cat/oocyte)[9][7]. Ecosystem assessments indicate that only limited habitat (tugai riparian zones in Kazakhstan/Uzbekistan) remains and that prey populations must be restored before any reintroduction[10][11]. Ethical and socio-political concerns are significant: experts question resource allocation and the welfare of engineered animals[12][13]. We conclude that while bioengineering tools make a Caspian-tiger-like animal theoretically possible, practical and ecological constraints make true revival unlikely in the near term. Future efforts might focus first on habitat restoration and conservation of Siberian tigers, alongside continued development of genome engineering and cloning methods.*

INTRODUCTION

The Caspian tiger (*Panthera tigris virgata*), also called the Turanian tiger, was a large feline inhabiting riverine forests and floodplains of Central Asia from Eastern Turkey through Iran, the Caucasus, and into Kazakhstan and western China[14]. Individuals reached up to about 530 pounds and 10 feet in length[4]. The subspecies was classified based on its distinct geographic range and morphology. Over the 20th century, Caspian tiger populations plummeted due to loss of riparian habitat (largely drained for Soviet irrigation and agriculture) and systematic killing[1][2]. By the 1950s the international community considered the Caspian tiger extinct[15], although isolated reports suggest a few may have persisted into the 1970s or even 1990s in remote areas of Turkey[16]. In any case, no confirmed wild Caspian tiger sightings have been recorded in decades.

Recent genetic studies have fundamentally changed our understanding of the Caspian tiger. Ancient-DNA analysis of museum specimens shows that Caspian and Siberian (Amur) tigers share a virtually identical mitochondrial genome, differing by only a single nucleotide[3]. Phylogeographic evidence suggests that a common *Panthera tigris* population colonized Central Asia and later Siberia via the Gansu Corridor about 10,000 years ago[3]. In practical terms, the living Amur tiger (*P. t. altaica*) is the closest extant relative to the Caspian tiger – so close that Driscoll et al. (2009) argue Amur tigers could serve as genetic surrogates if Caspian-range tigers were ever to be reintroduced[3]. Indeed, Luo et al. (2012) note that genetic difference between Amur and Caspian tigers is “negligible” and propose restoring tigers to the Caspian basin using Amur-stock animals[5].

In parallel to these findings, advances in biotechnology have raised the prospect of “de-extinction” – using genetic engineering to recreate extinct species or at least phenotypic analogs. Landmark projects include attempts to bring back the passenger pigeon, woolly mammoth, and Tasmanian tiger (thylacine) using CRISPR and cloning approaches (e.g., Church’s mammoth effort and Colossal Biosciences’ thylacine project)[6][7]. Recently, Colossal Biosciences reported success in creating live “dire wolf” pups by editing gray wolf DNA with ancient dire wolf sequences and using a domestic dog surrogate[8]. These developments have fueled interest in whether similar methods could be applied to the Caspian tiger.

This paper examines the feasibility of reviving the Caspian tiger via modern bioengineering. We review the tiger’s history and extinction causes, its genetic relationship to Siberian tigers, and the relevant biotechnologies: CRISPR/Cas9 gene editing, stem cell and embryo technologies, and interspecies nuclear transfer (cloning). We analyze biological challenges (DNA quality, phenotypic traits, reproductive biology), ecological considerations (remaining habitat and prey, ecosystem roles), technological constraints (current cloning success rates, surrogate compatibility), and ethical issues (resource priorities, animal welfare, ecosystem impacts). We then assess the potential ecological

impact of a restored tiger population in Uzbekistan and Central Asia. Our goal is to provide a comprehensive, evidence-based assessment of whether and how “bioengineering” could bring the Caspian tiger back to life, and what implications that would entail.




Fig. 1. Antique photograph of a captive Caspian tiger (*Panthera tigris virgata*). This extinct Central Asian subspecies was one of the largest tiger forms (up to ~530 lbs)[4]. Genetic data show Caspian tigers were essentially identical to Siberian (Amur) tigers[3].

Methods

This study synthesizes findings from peer-reviewed journals, conference proceedings, technical reports, and news accounts related to tiger conservation, de-extinction, and genome engineering. We identified key literature on Caspian tiger biology and extinction history[1][2], tiger genomics[3][5], CRISPR and genetic rescue technologies[6][7], and prior interspecies cloning attempts[12][7]. Sources included genetics studies (e.g. Driscoll et al. 2009, Chestin et al. 2017), high-profile de-extinction project reports (Shapiro 2015; Evans 2022), conservation organization analyses (WWF feasibility reports), and recent media on gene-edited animals (e.g. Kazinform 2025). When possible we extracted quantitative data on habitat size, tiger population models, genetic differences, and cloning outcomes. Technical details of CRISPR/Cas9 editing and somatic cell nuclear transfer (SCNT) were reviewed from molecular biology literature. We also reviewed ecological studies of Central Asian habitats to estimate available tiger range and prey base[17][11]. Ethical and socioeconomic commentary was gathered from opinion pieces and analyses where relevant (e.g. Bethge 2012; Evans 2022). All information is compiled to address the central question: can modern biotechnology realistically recreate a Caspian tiger and reintroduce it into the wild, and at what cost or impact?

RESULTS

Genetics and Historical Context

The Caspian tiger historically occupied scattered riparian forests (tugai vegetation) along Central Asian river systems[14]. By the early 1900s its range totaled on the order of 800,000–900,000 km²[18], but these were fragmented patches. Due to intensive hunting (including Soviet military bounty-hunts into the 1930s[2]) and habitat loss, Caspian tigers vanished rapidly. Soviet-era irrigation projects destroyed much of the tugai habitat[1], and prey species (Bukhara deer, wild boar, saiga) were decimated. By mid-century the subspecies was gone in the wild[15]. A recent analysis even suggests locals saw tigers in southeastern Turkey into the 1990s, implying some individuals survived decades longer than thought[16], but these claims remain unconfirmed.

Molecular studies have provided crucial insights. Driscoll et al. (2009) extracted mitochondrial DNA from museum specimens of Caspian tigers and found a nearly identical haplotype to that of modern Amur (Siberian) tigers[3]. The Caspian tiger's major mtDNA haplotype differed by only one nucleotide from the single type found in all living Amur tigers[3]. Phylogeographic analyses indicate that an ancestral *Panthera tigris* population migrated into Central Asia (~10,000 years ago) and later gave rise to the Amur tiger via eastward expansion[3]. Thus, genetically and evolutionarily, Caspian and Siberian tigers form a single clade. The implications are significant: living Amur tiger genomes effectively contain the Caspian tiger genome, minus any very rare mutations. Luo et al. (2012) concur that the difference between the two is “negligible” and propose using Amur tigers or their hybrids for reintroduction in the former Caspian range[5].

The limited genetic divergence simplifies the concept of bioengineering. Rather than constructing a wholly novel genome, one could start with a Siberian tiger genetic template and insert any Caspian-specific sequences identified from ancient DNA. Given that Caspian tigers may not have had unique traits beyond geographic population differences, the number of required gene edits might be small. However, most genetic differences likely reside outside mitochondrial DNA (in nuclear or regulatory genes), and comprehensive Caspian genome sequencing from degraded samples remains a challenge. Recent work on other extinct species underscores this point: Shapiro (2015) notes that resurrecting an extinct genome requires sequencing and comparing it to a living relative, then editing in targeted differences[19]. For species without living tissues, genome reconstruction from fossils is the only path. In principle, technology for extracting ancient DNA and assembling high-quality genomes has advanced greatly (now available for mammoths and passenger pigeons[19]), but it is imperfect: DNA damage and fragmentation in old specimens may leave gaps. Until a near-complete Caspian tiger genome is assembled, any bioengineering plan would rely on partial reconstructions and analogies with the Siberian tiger.

CRISPR and Genome Editing Techniques

CRISPR/Cas9 gene editing offers precise tools to rewrite DNA. In the context of de-extinction, scientists envision editing the genome of a close relative (or stem cell) to introduce traits of the extinct species. Shapiro (2015) describes the case of George Church's team, which in 2015 used CRISPR to replace 14 loci in an Asian elephant genome with the corresponding woolly mammoth sequences[6]. This involved identifying genes related to cold adaptation (fat insulation, hemoglobin, hair) and editing them in elephant cells. Shapiro notes that while fully recreating a mammoth genome would require ~1.5 million edits, focusing on key phenotypic genes could achieve an "Arctic elephant" phenotype[20]. Similarly, for Caspian tigers the number of meaningful edits could be far smaller, given the close relation. One could target genes involved in fur density, size, or adaptation to central Asian climates, if such differences exist.

In practice, CRISPR editing would proceed in cultured cells. One approach is to derive induced pluripotent stem cells (iPSCs) from a Siberian tiger fibroblast and use CRISPR to insert Caspian tiger alleles into those cells[21][7]. The engineered cells could be tested in vitro for expression and function. Alternatively, viral vectors or injected CRISPR complexes might be used directly in developing embryos (though this has lower efficiency). Shapiro emphasizes that as editing capability grows, we may be able to make extensive genomic changes even before fully understanding each gene's function[22]. For now, model organisms and cell assays would guide the process.

Once a suitably edited cell line is obtained, one would attempt to create an embryo. In mammals, the primary method is somatic cell nuclear transfer (SCNT), where the nucleus of an edited cell is transferred into an enucleated egg. If Caspian tigers were available, one might transfer a Caspian nucleus into a Siberian tiger egg; but with Caspians extinct, a Siberian (or domestic cat) oocyte would serve. The resulting embryo would carry the edited, Caspianized genome. Shapiro (2015) notes that SCNT has succeeded across many mammals and even in at least one extinct example: researchers cloned an extinct Pyrenean ibex (*bucardo*) in 2009[19]. That newborn ibex (born from a goat egg and mother) survived only minutes, but it demonstrated the principle of interspecies cloning (distantly related domestic goat surrogates). However, as Bethge (2012) and others report, cloning success rates are low and many interspecies embryos die prenatally or soon after birth[23]. A critical factor is genomic imprinting and epigenetics: foreign cytoplasm may mis-regulate transplanted nuclei. Shapiro (2015) emphasizes that if no suitable surrogate exists or if cloning fails, alternate methods (like germline genome editing in eggs/sperm) or artificial wombs might be needed[9]. For tigers, an adult cow or even domestic cat has successfully carried feline clones (tigers and leopards)[24], but a domestic animal is likely too distant for a tiger. A more plausible surrogate is a living Siberian tiger female; research has established Siberian tiger fibroblast cultures for this purpose[25]. Interspecies NT into a Siberian tiger egg is technically feasible in principle, though it may require optimization of

cell cycle stages and activation protocols.

Cloning and Hybridization Case Studies

To gauge the practicality of cloning an extinct felid, we examine analogous experiments. Bethge (2012) reports that interspecies cloning has been attempted for many endangered mammals: wildcats, lynxes, wolves, pandas, etc[24]. At the Audubon Center, for example, cloning of African wildcats and a black-footed cat using domestic cat eggs succeeded, producing live kittens[26]. These were not extinct subspecies but demonstrated that a domestic surrogate can gestate wild felid embryos. In the tiger domain, partial progress has been made: Song et al. (2007) cultured Siberian tiger fibroblasts specifically for nuclear transfer[25], providing donor cells with normal karyotypes. While no fully cloned tiger has been reported yet, these cell lines could serve as nuclei for SCNT.

A striking recent success with synthetic genomes was reported by Colossal Biosciences (2025) for the dire wolf. In this case, scientists identified dire wolf-specific gene variants (e.g. for coat and size) and used CRISPR to edit gray wolf cells. These gene-edited cells were then cloned (via SCNT) and implanted into domestic dog surrogates[8]. The result was three hybrid pups (“modern dire wolves”) that resemble the extinct *Aenocyon dirus*[8]. Two male pups were born in late 2024 and a female in early 2025[27]. The process involved transferring “healthy developing embryos...into surrogates for interspecies gestation”[28]. This real-world achievement shows that (1) CRISPR can incorporate extinct genes into a living relative, (2) SCNT can create embryos from those cells, and (3) surrogates can carry them to term. Importantly, it also highlights challenges: intensive monitoring was needed, and some experts question whether such hybrids can function ecologically[29].



Fig. 2. Dire wolf (*Aenocyon dirus*) pups born from CRISPR-edited gray wolf cells. Colossal

Biosciences introduced extinct wolf genes into living wolf cells and then cloned and carried the embryos in domestic dogs[8]. This demonstrates that gene-editing plus interspecies cloning can produce live animals with some extinct-animal traits.

The thylacine (Tasmanian tiger) project provides another instructive example of a marsupial de-extinction effort. Colossal Biosciences and University of Melbourne researchers plan to resurrect a thylacine-like animal by editing the genome of a related marsupial (fat-tailed dunnart) using CRISPR[7]. They have reprogrammed dunnart cells into stem cells and aim to insert thylacine sequences to create an embryo[7]. Because marsupial reproduction differs (tiny pouch young, short pregnancy), they are also developing artificial womb/pouch technologies[30]. While not a felid case, this project illustrates the workflow: sequence the extinct genome (the thylacine lab has >96% of it), identify target edits, reprogram relative's cells to iPSCs, apply CRISPR, and attempt to produce an embryo and surrogate birth[7][31]. The researchers note that gestation for thylacines would be much shorter than elephants, simplifying some aspects[31]. The thylacine effort underscores that emerging gene-editing methods can potentially "recreate" key extinct-trait combinations in a living proxy, but also that many steps (completing the genome, solving developmental challenges) remain.

In sum, published cases show that CRISPR-mediated de-extinction is no longer purely hypothetical. Mammalian cells can be edited with multiple extinct-species genes (Church's mammoth edits[6], WolfCorp's dire wolf[8]) and embryos brought to term via cloning. No cloned animal is a perfect genetic copy of its extinct progenitor, but it may express crucial traits. If the technological pipeline – ancient DNA recovery, genome assembly, targeted editing, SCNT, surrogate gestation – is successful, creating an animal very similar to the Caspian tiger is theoretically possible. However, the success rates of cloning in general remain low. Bethge (2012) notes that many cloned embryos die in utero or soon after birth[32]. The cloned Pyrenean ibex lived only 10 minutes, exemplifying high mortality[33]. Even the Dire Wolf project used modern grey wolves and dogs, whose genomes are well-known, whereas editing a tiger genome might encounter unforeseen viability issues.

Habitat and Ecosystem Feasibility

Apart from genetics, any revival must consider whether there is a suitable environment for Caspian tigers today. Luo et al. (2012) and subsequent studies have examined Central Asian habitat. Large areas of potential habitat remain: Luo et al. identified over 1 million km² of former tiger range that could be suitable if restoration measures were taken[34]. Chestin et al. (2017) used remote sensing and population models to narrow this to a few patches. They found that only two key areas (totaling <20,000 km²) are currently viable for a tiger population[35]. The most promising is the Ili River delta and southern Lake Balkhash region in eastern Kazakhstan (~7,000 km² of high-quality riparian forest)[11]. In this region, models suggest that a population of ~40–50 tigers could grow to 64–98 in 50 years, provided prey and water are plentiful[11]. These

estimates assume careful habitat management: maintaining Ili River flow, preventing floodplain fires, and bolstering prey (Bukhara deer, wild boar, roe deer)[11][10].

Extensive surveys in Uzbekistan's Amu Darya Delta (Tugai forests) have identified the Akpetki–Jiltyrbas area (~1 million ha) as a candidate site[17]. However, experts emphasize that it is not yet ready for tigers. WWF consultants recommend first securing protected-area status, restoring water flow, and rebuilding prey stocks[17][10]. For example, current ungulate densities are far below historical levels, and key species like the kulan (Asiatic wild ass) and Bukhara deer need reintroduction[10]. Livestock grazing and oil/gas development must be strictly managed or removed to prevent human–tiger conflict[17][10]. In short, the ecological infrastructure for tigers (dense prey, cover, water) must be deliberately reconstructed. This aligns with Chestin et al.'s conclusion that tiger re-establishment is feasible “if...prey population restoration precedes tiger introduction”[36].

Community attitudes are also a factor. In Kazakhstan's Ili-Balkhash region, the local Auyldastar community has been engaged in preparation: planting native trees, building fire brigades, and learning about tigers[37][38]. Initially there was hesitation, but after six years of outreach, local leaders express pride in the project[39]. The Siberian tigers being introduced (named Bodhana and Kuma) are kept first in large enclosures. If they successfully breed, their cubs are intended for release into the wild[40]. This ongoing real-world trial (launched 2022–2025) will provide concrete data on ecosystem carrying capacity and socio-ecological dynamics. It illustrates that even without de-extinction, reintroducing close relatives into the Caspian range is possible with long-term commitment[41][39].

From the ecosystem perspective, tigers are apex predators whose presence or absence can have cascading effects. Restoration of wolves in Yellowstone, for example, led to significant trophic changes. Experts suggest that if tigers reach viable densities, they could help control overabundant herbivores (as the thylacine project anticipates for wallabies and kangaroos[42]). In Central Asia, tigers would prey on wild boars, deer, gazelles, and perhaps reduce pressure on vegetation. Conversely, introducing a large predator carries risks: livestock depredation could provoke human backlash. Plans therefore typically include livestock compensation programs and strict anti-poaching enforcement[10].

Technological and Biological Challenges

Genome Reconstruction: A revived Caspian tiger genome must account for all missing DNA, not just a few traits. The Church lab's mammoth project highlights that even when high-coverage genomes are available, tens of thousands of edits might be needed to approximate the extinct species[20]. If Caspian tigers diverged ~10,000 years ago, their genomes could differ at many loci. We lack published estimates of nuclear divergence between Caspian and Siberian tigers; mitochondrial evidence shows near identity, but the

nuclear genome could harbor unique alleles affecting coat pattern, disease resistance, or other physiology. Ancient DNA from museum skins can help identify fixed differences, but quality issues (fragmentation, contamination) may leave gaps. As Shapiro warns, “the number of edits can be minimized by replacing large pieces of the genome in a single edit or by focusing on phenotypic genes”[20], but this requires knowing which genes matter. Without complete knowledge of gene–trait links, CRISPR edits may miss key variants. There is also risk of off-target effects or incomplete editing.

Cloning and Embryo Viability: Even if a designer Caspian genome is assembled, producing a healthy embryo is nontrivial. Shapiro (2015) notes that cloning elephants has not yet been accomplished[9]; similarly, no successful tiger SCNT with a foreign nucleus has been reported publicly. The only tiger embryos ever gestated outside a tiger surrogate were in vitro, and any live births would be unprecedented. Interspecies SCNT often faces epigenetic barriers: mismatch between nucleus and cytoplasm can cause developmental arrest or abnormalities (Bethge 2012). Indeed, many animal cloning attempts yield stillbirths or postnatal death[43]. Even Colossal’s dire wolf pups required careful optimization and currently live under intensive care. If a Caspian tiger embryo were created, a suitable surrogate would likely be an Amur tiger. However, Siberian tigers themselves are endangered (~500 in captivity worldwide) and captive breeding programs cannot easily spare females for experimentation. The other option would be to use a domestic surrogate (e.g. cat or dog), but such distant species may fail to support tiger embryo development. Shapiro suggests artificial wombs as a future solution for intractable cases[9], but these are still experimental for mammals.

Population Genetics: Another biological hurdle is genetic diversity. Any de-extinct Caspian tiger population would start from a very small founder pool (perhaps one or a few reconstructed genomes), leading to inbreeding. Even the dire wolves created are essentially three individuals from a single plan. In contrast, natural populations benefit from many founders over time, preserving genetic health. To avoid severe inbreeding depression, one would need multiple independent cell lines or gametes reflecting any known genetic variation among Caspian tigers (which were already somewhat bottlenecked before extinction). The 2009 study of 20 Caspian specimens found “genetic depletion” characteristic of Amur tigers, indicating low diversity predating human effects[44]. Reviving a genetically diverse population would likely require mixing multiple edited genomes or repeatedly introducing gene-edited individuals (akin to a breeding program) – a complex, ongoing process.

Health and Adaptation: Even if live Caspian-like tigers were created, their health and viability are uncertain. Cloned animals often exhibit higher rates of developmental defects, immune issues, and abnormal aging[32]. For example, the cloned bucardo died of lung problems after birth[33]. The same risk could affect cloned or gene-edited tigers. Furthermore, Caspian tigers were adapted to a specific environment (riparian Central

Asia). Modern surrogates (Siberian tigers) and potential offspring may have phenotypes optimized for Siberian forests, not arid steppes. If gene edits focus only on visible traits (coat, size), subtler physiological adaptations (salt tolerance, fat storage) might be missed, reducing fitness. Thus, a revived Caspian tiger might be only an approximation.

Ethical, Financial, and Social Considerations

Many scientists caution against de-extinction as a conservation panacea. Critics argue it may divert resources from saving extant species and habitats. As Bethge (2012) reports, some experts contend that de-extinction “trivializes extinction” and that funds would be better spent on habitat protection[12]. Indeed, the resources needed for gene-editing projects are immense; Colossal has raised hundreds of millions for mammoth and thylacine work. For a Caspian tiger project, similar investment would be required for research facilities, sequencing, editing, and long-term captive breeding. Meanwhile, Asiatic tiger subspecies (including Amur tigers) remain critically endangered, and Central Asian habitats still lack basic protections.

Animal welfare is another concern. Cloned animals often suffer abnormalities and large-scale cloning of embryos could cause suffering (high failure rates, stillbirths). Some ethicists question whether creating a novel, likely-hybrid animal in artificial conditions is morally acceptable, especially if the end goal is release into uncertain ecosystems. Preston (2025) summed up a common skepticism: “It is hard to imagine dire wolves ever being released and taking up an ecological role.”[13] If even supporters doubt releasing a hybrid wolf, similar doubts apply to tigers whose historical niche no longer exists as it once did.

On the other hand, proponents argue that technology itself can aid conservation. The thylacine researchers note that developing marsupial reproductive techniques (e.g., artificial pouch, milk formula) may benefit living endangered marsupials[31]. Similarly, somatic cell lines or frozen DNA from tigers could serve as genetic “insurance” for future use, even if full revival is not done. Furthermore, a restored tiger could catalyze ecosystem protection; Chestin et al. observe that tiger reintroduction programs often lead to new protected areas and funding for biodiversity[36]. A flagship charismatic species can galvanize public support. In Kazakhstan, the government and WWF have partnered since 2011 to restore landscapes (Balkhash-Balkans River basins) in anticipation of tigers[45], showing a political will that could extend to Uzbekistan.

Social factors are crucial. Reintroducing any tiger (genetically revived or not) requires community consent. The Kazakhstan project demonstrates that local attitudes can shift with engagement and economic benefits (ecotourism, grants). Conversely, if communities fear livestock losses, conflicts may undermine conservation. The WWF feasibility report recommends compensation schemes and awareness campaigns[10]. In Uzbekistan and elsewhere, similar groundwork would be essential.